

Pertussis Disease (Whooping Cough)

What is pertussis disease?

- Pertussis, also known as whooping cough, is a highly contagious bacteria respiratory disease.
- Pertussis is known for uncontrollable, violent coughing which often makes it hard to breathe.
- After cough fits, someone with pertussis often needs to take deep breaths, which result in a “whooping” sound.
- Pertussis can affect people of all ages, but can be very serious, even deadly, for babies less than a year old.
- Pertussis is a highly contagious disease and can affect up to 80% household contacts.

What are the symptoms?

- Symptoms of pertussis usually develop within 5 to 10 days after you are exposed, but may not develop for as long as 3 weeks.
- First week or two, early symptoms may include:
 - Subtle onset of runny nose.
 - Low grade fever.
 - Sneezing.
 - Mild, occasional cough.
 - Apnea or stop breathing when asleep.
- After 1 to 2 weeks, symptoms may include:
 - Fits of many, rapid coughs followed by a high-pitched “whoop” sound..
 - Throwing up during or after coughing fits.
 - Very tired after coughing fits.

How does pertussis spread?

- Pertussis is spread from person-to person.
 - Coughing and sneezing.
 - Spend a lot of time near one another and share breathing space.
 - Some babies get pertussis from older siblings, parents, or caregivers who might not even know they have the disease.
- The best prevention for pertussis is vaccine, but no vaccine is 100% effective.

Where can I find more information?

- Ask your doctor, or your local county health department.
- Email the Alabama Department of Public Health, Immunization Division, at immunization@adph.state.al.us.
- Go to cdc.gov and type 'pertussis disease' in the SEARCH box.



Alabama Department of Public Health

Immunization Division, 201 Monroe St, Montgomery, AL 36104

1-800-469-4599 www.adph.org/imm 2/29/16

Pertussis Vaccine

Who should get pertussis vaccine?

- There are 2 vaccines which protect against pertussis. DTaP for children up to 6 years of age and Tdap for people 7 years of age and older.
- People who have not had been vaccinated and are in n contact with a baby.
- Babies beginning at 2 months of age, followed by a total of five doses through age 4-6 years.
- Adolescents aged 11 through 18 years should receive a single Tdap.
- Adults aged 19 through 64 years should receive a single Tdap.
- Pregnant women should receive a Tdap for each pregnancy.
- Anyone who has not received the full five doses of pertussis vaccine.
- Either if you are exposed to pertussis and/or have the disease, you should still receive vaccine containing pertussis.
- All adults are recommended one dose of Tdap as soon as possible, especially if pregnant or in close contact with infants.

What are the vaccine side effects and risks?

- Mild (common) problems after the vaccine may include:
 - Fever.
 - Pain, redness, and/or swelling where shot was given.
 - Local reactions such as pain, redness, or swelling.
 - Drowsiness, fretfulness.
 - Tiredness.
- Moderate (uncommon) problems after the vaccine may include:
 - Seizure (jerking or staring).
 - Non-stop crying, for 3 hours or more.
 - High fever, over 105°F.
- Severe (very rare) problems after the vaccine may include:
- Serious allergic reaction.

Where can I find more information?

- Ask your doctor, or your local county health department.
- Email the Alabama Department of Public Health, at immunization@adph.state.al.us.
- Go to cdc.gov and type 'pertussis vaccine' in the SEARCH box.



Alabama Department of Public Health

Immunization Division, 201 Monroe St, Montgomery, AL 36104

1-800-469-4599 www.adph.org/imm 2/29/16