

## Minimum Immunization Requirements for Those Entering a Child Care Facility or School in Illinois, Fall-2014

Vaccine Requirement	Child Care Facility, Preschool, Early Childhood Pre-Kindergarten Programs	Kindergarten through 12 <sup>th</sup> Grade		Minimum Intervals Allowed Between Doses and Other Options for Proof of Immunity
		First Entry into School (Kindergarten or First Grade)	Other Grades	
<b>DTP/DTaP/ or Tdap, Td</b> (Diphtheria, Tetanus, Pertussis)	Three doses by 1 year of age  One additional booster dose by 2 <sup>nd</sup> birthday	Four or more doses of DTP/DTaP with the last dose qualifying as a booster and received on or after the 4 <sup>th</sup> birthday	Three or more doses of DTP/DTaP or Td with the last dose qualifying as a booster and received on or after the 4 <sup>th</sup> birthday  <b>Beginning school year 2013-2014*: Students entering 6<sup>th</sup> thru 12<sup>th</sup> grades: 1 dose of Tdap</b>	Between series doses: 4 weeks (28 days)  Between series and booster: 6 months
<b>Polio</b>	Two doses by 1 year of age.  One additional dose by 2 <sup>nd</sup> birthday	Three or more doses of the same type of Polio vaccine with the last dose qualifying as a booster and received on or after the 4 <sup>th</sup> birthday. *If the series is given in any combination of polio vaccine types, 4 or more doses are required with the last being a booster on or after the 4 <sup>th</sup> birthday.	Three or more doses of Polio with the last dose qualifying as a booster and received on or after the 4 <sup>th</sup> birthday.  *If the series is given in any combination of polio vaccine types, 4 or more doses are required with the last being a booster on or after the 4 <sup>th</sup> birthday.	Minimum interval between series doses: 4 weeks (28 days)
<b>Measles</b>	One dose on or after the 1 <sup>st</sup> birthday but prior to 24 months of age	Two doses of Measles Vaccine, the 1 <sup>st</sup> dose must have been received on after the 1 <sup>st</sup> birthday and the second dose no less than 4 weeks (28 days) later.		<i>Laboratory evidence of measles immunity OR Certified physician verification* of measles disease by date of illness</i> <b>*Cases diagnosed after 7/1/02 must include lab evidence</b>
<b>Rubella</b>	One dose on or after the 1 <sup>st</sup> birthday but prior to 24 months of age	Two doses of Rubella Vaccine, the 1 <sup>st</sup> dose must have been received on after the 1 <sup>st</sup> birthday and the second dose no less than 4 weeks (28 days) later.		<i>Laboratory evidence of rubella immunity</i> <b>History of disease is not acceptable proof of immunity to rubella</b>
<b>Mumps</b>	One dose on or after the 1 <sup>st</sup> birthday but prior to 24 months of age	Two doses of Mumps Vaccine, the 1 <sup>st</sup> dose must have been received on after the 1 <sup>st</sup> birthday and the second dose no less than 4 weeks (28 days) later.		<i>Laboratory evidence of mumps immunity OR Certified physician verification of mumps disease by date of illness</i>
<b>Haemophilus influenzae type b</b>	Refer to Hib vaccination schedule for series Children 24-59 mos. without series must have one dose after 15 mos. of age	<b>Not required after the 5<sup>th</sup> birthday (60 months of age)</b>		Refer to Hib vaccination schedule
<b>Hepatitis B</b>	Three doses for all children 2 years of age or older  Third dose must have been administered on or after 6 months of age (168 days)	<b>No Requirements</b>		Minimum intervals between doses: 1 & 2 - at least 4 weeks (28 days) 2 & 3 - at least 2 months (56 days) 1 & 3 - at least 4 months (112 days) <i>Laboratory evidence of prior or current infection</i>
<b>Varicella</b>	One dose on or after the 1 <sup>st</sup> birthday	<b>Two doses of Varicella Vaccine</b> , the 1 <sup>st</sup> dose must have been received on after the 1 <sup>st</sup> birthday and the second dose no less than 4 weeks (28 days) later.		<i>Statement from physician or health care provider verifying disease history OR Laboratory evidence of varicella immunity</i>
		<b>Applies to Children entering grades 6 thru 12 for 2014-2015 school year</b>  Three doses administered at recommended intervals		<b>Applies to Students entering grades 1 thru 12</b> One dose of Varicella on or after the 1 <sup>st</sup> birthday  <b>Applies to Students entering 6<sup>th</sup> or 9<sup>th</sup> grade beginning 2014-2015 school year</b> Two doses of Varicella Vaccine, the 1 <sup>st</sup> dose must have been received on after the 1 <sup>st</sup> birthday and the second dose no less than 4 weeks (28 days) later  <i>Statement from physician or health care provider verifying disease history OR Laboratory evidence of varicella immunity</i>
<b>Invasive Pneumococcal Disease</b>	Refer to PCV vaccination schedule for series Children 24-59 mos. without series must have one dose	<b>Not required after the 5<sup>th</sup> birthday (60 months of age)</b>		Refer to PCV vaccination schedule

D-1

**Section 695.APPENDIX A Vaccination Schedule for Haemophilus influenzae type b Conjugate Vaccines (Hib)**

Vaccine	Age at 1 <sup>st</sup> of doses for dose (mos.)	Primary series	Booster	Total number series
HbOC/PRP-T HibTITER™	2-6	3 doses, 2mo. apart <sup>a</sup>	12-15 mo. <sup>bc</sup>	4
	7-11	2 doses, 2mo. apart <sup>a</sup>	12-18 mo. <sup>bc</sup>	3
	12-14	1 dose	15 mo. <sup>bc</sup>	2
ActHib <sup>e</sup> ™ OmniHib™ TETRAMUNE™	15-59	1 dose <sup>d</sup>	None	1
PRP-OMP PedvaxHIB™	2-6	2 doses, 2mo.apart <sup>a</sup>	12 mo. <sup>bc</sup>	3
	7-11	2 doses, 2mo.apart <sup>a</sup>	12-18 mo. <sup>bc</sup>	3
	12-14	1 dose	15 mo. <sup>bc</sup>	2
	15-59	1 dose <sup>d</sup>	None	1
PRP-D ProHIBIT™	15-59	1 dose <sup>cd</sup>	None	1

<sup>a</sup> Minimally acceptable interval between doses is one month.

<sup>b</sup> At least 2 months after previous dose.

<sup>c</sup> After the primary infant Hib vaccine series is completed, any of the licensed Hib conjugate vaccines may be used as a booster dose.

<sup>d</sup> Children 15-59 months of age should receive only a single dose of Hib vaccine.

<sup>e</sup> Reconstituted with DTP as a combined DTP/Hib vaccine

Note: A DTP/Hib combination vaccine can be used in place of HbOC or PRP-T

(Source: Amended at 26 Ill. Reg. 10792, effective July 1, 2002)

**Section 695.Appendix B Vaccination Schedule for Pneumococcal Conjugate Vaccines (PCV)**

<b>Age of Child (Months)</b>	<b>Vaccination History</b>	<b>Primary Series and Booster Intervals</b>	<b>Total Doses Required</b>
<b>2-6 minimum age of six weeks:</b>	0 doses	3 doses, 2 months apart; 4 <sup>th</sup> dose at age 12-15 months	4
	1 dose	2 doses, 2 months apart; 4 <sup>th</sup> dose at age 12-15 months	4
	2 doses	1 dose, 2 months after most recent dose; 4 <sup>th</sup> dose at age 12-15 months	4
<b>7-11</b>	0 doses	2 doses, 2 months apart; 3 <sup>rd</sup> dose at age 12 -15 months	3
	1 or 2 doses before age 7 months	1 dose, 2 months after most recent dose; 3 <sup>rd</sup> dose at 12 months -15 months and > 2 months after prior dose	3-4
<b>12-23</b>	0 doses	2 doses, $\geq 2$ months apart	2
	1 dose administered before age 12 months	2 doses, $\geq 2$ months apart	2
	1 dose administered on or after 12 months of age	1 dose $\geq 2$ months after most recent dose	2
	2 or 3 doses administered before age 12 months	1 dose, $\geq 2$ months after most recent dose	3-4
<b>24-59 Healthy Children</b>	Any incomplete schedule	1 dose, $\geq 2$ months after most recent dose	1
Children at High risk <sup>II</sup>	Any incomplete schedule	2 doses separated by 2 months	2