

These Immunizations are recommended for your child.

Child's Age	Vaccine and Dose	Protects Against
Birth 2 months (part of well-baby visit)	Hepatitis B Dose 1 of 3	Hepatitis B virus (chronic inflammation of the liver, life-long complications)
	DTaP Dose 1 of 5	Diphtheria, tetanus, and acellular pertussis (whooping cough)
	Hib Dose 1 of 4	Infections of the blood, brain, joints, or lungs (pneumonia)
	Polio (IPV) Dose 1 of 4	Polio
	Hepatitis B Dose 2 of 3	Hepatitis B virus (chronic inflammation of the liver, life-long complications)
	Pneumococcal conjugate (PCV13) Dose 1 of 4	Infections of the blood, brain, joints, or inner ears
	Rotavirus Dose 1 of 3	Rotavirus (severe diarrhea and vomiting)
4 months (part of well-baby visit)	DTaP Dose 2 of 5	Diphtheria, tetanus, and acellular pertussis (whooping cough)
	Hib Dose 2 of 4	Infections of the blood, brain, joints, or lungs (pneumonia)
	Polio (IPV) Dose 2 of 4	Polio
	Pneumococcal conjugate (PCV13) Dose 2 of 4	infections of the blood, brain, joints, or inner ears
	Rotavirus Dose 2 of 3	Rotavirus (severe diarrhea and vomiting)
6 months (part of well-baby visit)	DTaP Dose 3 of 5	Diphtheria, tetanus, and acellular pertussis (whooping cough)
	Hib Dose 3 of 4	Infections of the blood, brain, joints, or lungs (pneumonia)
	Pneumococcal conjugate (PCV13) Dose 3 of 4	Infections of the blood, brain, joints, or inner ears
	Rotavirus Dose 2 of 3 no later than 32 weeks	Rotavirus (severe diarrhea and vomiting)
6 to 18 months	Hepatitis B Dose 3 of 3	Hepatitis B (chronic inflammation of the liver, life-long complications)
	Polio (IPV) Dose 3 of 4	Polio

6 months through 18 years	<u>Influenza</u> 1 dose every year	Flu and complications
12 to 15 months	<u>Hib</u> Dose 4 of 4	Infections of the blood, brain, joints, or lungs (pneumonia)
	<u>Pneumococcal conjugate (PCV13)</u> Dose 4 of 4	Infections of the blood, brain, joints, or inner ears
	<u>MMR</u> Dose 1 of 2	Measles, mumps, and rubella (German measles)
	<u>Hepatitis A</u> Dose 1 of 2	Hepatitis A virus (chronic inflammation of the liver, life-long complications)
12 to 18 months	<u>Varicella</u> Dose 1 of 2	Chickenpox
15 to 18 months	<u>DTaP</u> Dose 4 of 5	<u>Diphtheria, tetanus, and acellular pertussis (whooping cough)</u>
	<u>Hepatitis A</u> Dose 2 of 2	Hepatitis A virus (chronic inflammation of the liver, life-long complications)
2 to 5 years <i>Catch-up immunization requirement, if previous 4 PCV's were not met.</i>	<u>Pneumococcal conjugate (PCV13)</u> Dose 1 of 2 if using catch-up schedule	Infections of the blood, brain, joints, or inner ears
5 years and older, through 18 years certain high-risk groups	<u>Pneumococcal conjugate (PCV13)</u> Dose 2 of 2, if needed (follows 3-5 years after Dose 1)	Infections of the blood, brain, joints, or inner ears
4 to 6 years	<u>DTaP</u> Dose 5 of 5	<u>Diphtheria, tetanus, and acellular pertussis (whooping cough)</u>
	<u>Polio (IPV)</u> Dose 4 of 4	Polio
	<u>MMR</u> Dose 2 of 2	Measles, mumps, and rubella (German measles)
	<u>Varicella</u> Dose 2 of 2	Chickenpox
11 to 12 years	Td booster, 1 dose every 10 years	<u>Tetanus</u> and <u>diphtheria</u>
	Tdap (single dose in place of Td booster)	<u>Tetanus, diphtheria</u> and <u>acellular pertussis (whooping cough)</u>
	<u>HPV</u> (3 doses total)	Human Papilloma Virus
	<u>MCV4</u>	Meningococcal conjugate (prevents meningitis)
Additional Vaccines	Consult your child's physician or healthcare professional about catch-up vaccinations and vaccines for travelers .	

