

## Recommendations for Hydration to Prevent Heat Illness

### Types Of Sports Drinks

➤ **Fluid Replacers**

Examples: Water, Gatorade, 10K, Quickkick, Max

These drinks are absorbed as quickly as water and typically are used for activities lasting less than two hours.

➤ **Carbohydrate loaders**

Examples: Gatorlode, Exceed High, Carboplex

These drinks replace more muscle glycogen to enhance greater endurance.

They should be used after ultra-endurance events to increase muscle glycogen resynthesis after exercise.

➤ **Nutrition Supplements**

Examples: Gatorpro, Exceed Sports, Ultra Energy

These supplements are fortified with vitamins and minerals and they help athletes maintain a balanced diet.


They can be used as a meal replacement supplement for athletes who wish to skip a high fat meal, or as extra calories for athletes who wish to gain weight.

### What Not To Drink

- Drinks with Carbohydrate (CHO) concentrations of greater than eight percent should be avoided.
- Fruit juices, CHO gels, sodas, and sports drinks that have a CHO greater than six to eight percent are not recommended during exercise as sole beverages.
- Beverages containing caffeine, alcohol, and carbonation are not to be used because of the high risk of dehydration associated with excess urine production, or decreased voluntary fluid intake.

## Hydration Tips And Fluid Guidelines

- Drink according to a schedule based on individual fluid needs.
- Drink before, during and after practices and games.
- Drink 17-20 ounces of water or sports drinks with six to eight percent CHO, two to three hours before exercise.
- Drink another 7-10 ounces of water or sport drink 10 to 20 minutes before exercise.
- Drink early - By the time you're thirsty, you're already dehydrated.
- In general, every 10-20 minutes drink at least 7-10 ounces of water or sports drink to maintain hydration, and remember to drink beyond your thirst.
- Drink fluids based on the amount of sweat and urine loss.
- Within two hours, drink enough to replace any weight loss from exercise.
- Drink approximately 20-24 ounces of sports drink per pound of weight loss.
- Dehydration usually occurs with a weight loss of two percent of body weight or more.



## What To Drink During Exercise

- If exercise lasts more than 45-50 minutes or is intense, a sports drink should be provided during the session.
- The carbohydrate concentration in the ideal fluid replacement solution should be in the range of six to eight percent CHO.
- During events when a high rate of fluid intake is necessary to sustain hydration, sports drinks with less than seven percent CHO should be used to optimize fluid delivery. These sports drinks have a faster gastric emptying rate and thus aid in hydration.
- Sports drinks with a CHO content of 10 percent have a slow gastric emptying rate and contribute to dehydration and should be avoided during exercise.
- Fluids with salt (sodium chloride) are beneficial to increasing thirst and voluntary fluid intake as well as offsetting the amount of fluid lost with sweat.
- Salt should never be added to drinks, and salt tablets should be avoided.
- Cool beverages at temperatures between 50 to 59 degrees Fahrenheit are recommended for best results with fluid replacement.